

E 20 2060

PL-TR-93-2236

AD-A282 009



**EFFECT OF COULOMB COLLISIONS, LANDAU
DAMPING AND PARTICLE TRAPPING ON
LINEAR AND NONLINEAR ION ACOUSTIC WAVES**

**Chung Chan
Zhijiang Jin**

**DTIC
ELECTE
JUL 06 1994
S F D**

**Northeastern University
Department of Electrical Engineering
Boston, MA 02115**

December 30, 1993

Scientific Report No. 1

16 p 94-20520

Approved for public release; distribution unlimited

DTIC QUALITY INSPECTED 3



**PHILLIPS LABORATORY
Directorate of Geophysics
AIR FORCE MATERIEL COMMAND
HANSCOM AIR FORCE BASE, MA 01731-3010**

94 7 5 188

"This technical report has been reviewed and is approved for publication"

Bamandas Basu

(Signature)

BAMANDAS BASU

Contract Manager

David N. Anderson

(Signature)

DAVID N. ANDERSON

Branch Chief

William K. Vickery

(Signature)

WILLIAM K. VICKERY

Division Director

This report has been reviewed by the ESC Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center (DTIC). All others should apply to the National Technical Information Service (NTIS).

If your address has changed, if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify PL/TSI, 29 Randolph Road, Hanscom AFB, MA 01731-3010. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 30 December 1993	3. REPORT TYPE AND DATES COVERED Scientific No. 1
4. TITLE AND SUBTITLE Effect of Coulomb Collisions, Landau Damping and Particle Trapping on Linear and Nonlinear Ion Acoustic Waves			5. FUNDING NUMBERS PE 61102F PR 2310 TA G6 WU EB Contract F19628-92-K-0011
6. AUTHOR(S) Chung Chan Zhijiang Jin			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Northeastern University Department of Electrical Engineering Boston, MA 02115			8. PERFORMING ORGANIZATION REPORT NUMBER
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Phillips Laboratory 29 Randolph Road Hanscom AFB, MA 01731-3010 Contract Manager: Bamandas Basu/GPIM			10. SPONSORING/MONITORING AGENCY REPORT NUMBER PL-TR-93-2236
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 words) We have performed a series of experiments to study the damping effect of the ion acoustic waves. The experimental results have been analyzed both qualitatively and quantitatively based on the theory by Basu and Jasperse ^[1] . The analysis shows that the theoretical and experimental results agree very well under certain conditions.			
14. SUBJECT TERMS Ion acoustic waves Collision Damping			15. NUMBER OF PAGES 16 16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT SAR

CONTENTS

1. INTRODUCTION	1
2. PROGRESS SUMMARY	1
3. PROGRESS REPORT	1
3.1. Experimental Setup.....	1
3.2. Theory and Experiment on the Ion Acoustic Waves	2
4. PROPOSED RESEARCH.....	4
References.....	5

ILLUSTRATIONS

1. Schematic of the Plasma Chamber for Measuring Ion Acoustic Waves.....	6
2. The I-V Curve Taken From Langmuir Probe.	7
3. The Waves Measured in Different Frequencies.....	8
4. Normalized Wave Number and Damping Rate vs Ratio of Ion to Electron Temperatures.	10
5. The Wave Number and Damping Rate vs Frequency.....	11
6. (a). Wave Number vs Frequency; (b). Damping Rate vs Frequency	12

TABLE

1. Experimental and Theoretical Wave Parameters.....	6
--	---

NTIS CRA&I	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

1. INTRODUCTION

Our project is to investigate experimentally the effect of Landau damping and particle collisions on the ion acoustic waves. For the collisionless current-free plasma, linear Vlasov theory predicts that ion acoustic waves are Landau damped and plasma is stable. For the weakly collisional plasma, a recent theory^[1] indicates that electron-ion collisions have an undamping effect on the ion acoustic waves. If this undamping effect exceeds other damping effects, the plasma will be unstable. We have performed a series of experiments to study the damping effect of the ion acoustic waves. The experimental results are also explained qualitatively and quantitatively based on the reference [1].

2. PROGRESS SUMMARY

A series of experiments have been performed during last fiscal year. The experimental results have been analyzed both qualitatively and quantitatively based on the theory by *Basu and Jasperse*^[1]. The analysis shows that the theoretical and experimental results agree very well under certain conditions.

3. PROGRESS REPORT

3.1. Experimental Setup

The experiments were performed in a double plasma chamber as shown in Figure 1. The plasma was produced by impact ionization from hot filament cathodes at the top and bottom ends. The plasma potential, the electron temperature and density were measured with a movable Langmuir probe and the ion temperature was measured with a gridded energy analyzer. A program has been written to extract the plasma parameters from the I-V curve of Langmuir probe automatically. Figure 2 shows a set of measured I-V curves and also the extracted plasma parameter values. Usually we measured one I-V curve for

every 5cm and totally four curves (at $x=5\text{cm}$, 10cm , 15cm , and 20cm). The typical plasma parameters used in the experiment are as follows: electron temperature $T_e=1\sim 2\text{eV}$, ion temperature $T_i=0.1\sim 0.2T_e$, electron density $n_e=10^8\sim 10^{10}\text{cm}^{-3}$, total argon gas pressure $p\approx 10^{-4}\text{ Torr}$, and base pressure $p_b\approx 10^{-6}\text{ Torr}$.

Waves were launched externally with a signal grid (30cm in diameter, 34cm away from the bottom end) and were detected by the movable Langmuir probe. An example of the measured waves is shown in Figure 3. Notice that for each frequency we took two sets of data.

3.2. Theory and Experiment on the Ion Acoustic Waves

For the collisional plasma, *Basu and Jasperse*^[1] gave the following approximate formulas for the ion acoustic waves:

$$k_r \lambda_e = \frac{\omega}{\omega_{pi}} \left(1 + \frac{3}{\theta}\right)^{-1/2} \quad (1a)$$

$$(k_i \lambda_e)^{\text{Landau}} = \frac{\omega}{\omega_{pi}} F(\theta) \quad (1b)$$

$$(k_i \lambda_e)^{\text{ei}} = -\frac{1}{4} \frac{v_{ei}}{\omega_{pi}} \left(1 + \frac{3}{\theta}\right)^{-1} \quad (1c)$$

$$(k_i \lambda_e)^{\text{in}} = \frac{1}{2} \frac{v_{in}}{\omega_{pi}} \left(1 + \frac{3}{\theta}\right)^{-2} \quad (1d)$$

where

$$F(\theta) = \left(1 + \frac{3}{\theta}\right)^{-1} \left(\frac{\pi}{8}\right)^{1/2} \left[\left(\frac{m_e}{m_i}\right)^{1/2} + \theta^{3/2} \exp\left(-\frac{\theta}{2} - \frac{3}{2}\right) \right], \quad \theta = \frac{T_e}{T_i},$$

v_{ei} and v_{in} are electron-ion and ion-neutral collision frequency, ω_{pi} is the plasma frequency of ion, λ_e is the Debye length of electron, and perturbations are taken as $e^{i(kx-\omega t)}$. The wave number and the Landau damping can also be approximated from the collisionless kinetic equation which is given as:

$$\varepsilon(\omega, k) = 1 - \frac{1}{2k^2\lambda_e^2} Z'\left(\frac{\omega}{ka_e}\right) - \frac{1}{2k^2\lambda_i^2} Z'\left(\frac{\omega}{ka_i}\right) = 0 \quad (2)$$

where λ_i , a_e , a_i and Z' denote the Debye length of ions, the thermal velocities of electron and ion, and the derivative of the plasma dispersion function respectively. In Figure 4, both results are shown. The solid line in the figure is calculated from Formula (1a) and (1b) and the dashed line is calculated from Equation (2). It is obvious that the formulas give good approximate values when T_i/T_e is small, but there are large differences between the approximate formula and the solutions from the kinetic equation when T_i/T_e is large. Since we usually have $T_i/T_e \geq 0.1$ in our experiments, Equation (2) was used to calculate the wave number and the Landau damping rate.

For the wave data shown in Figure 3, Figure 5 shows the results of wave number and damping rate obtained from the curve fitting. From the theory we know that the wave number and the damping rate are approximately a linear function of the frequency. By line fitting the data in Figure 5 we derive the wave parameters, which are shown in Table 1. Note that when the frequency is less than 70kHz the experimental damping rate does not follow the theoretical prediction. Two reasons may contribute to this disagreement: we note that this occurs when the wave frequencies are near or less than the electron-ion collision frequency which is about 61kHz and the wavelength is near the size of our chamber. We can see that the theoretical and experimental values are very close to each other. The parameters we used are: $n_e \approx 4.62 \times 10^{12} \text{ cm}^{-3}$, $n_i \approx 1.27 \times 10^9 \text{ cm}^{-3}$, $T_e \approx 0.84 \text{ eV}$, and $T_i \approx 0.16 \text{ eV}$.

We also measured the ion acoustic waves in different plasma densities. Figure 6 show the results of wave numbers and damping rates in three different density cases. From Figure 6a, we find that the wave speeds for three density cases are quite close. From Figure 6b, we find that the damping rate decreases as the density increases just as the theory predicts. However, in the high density region, we find that the damping rate becomes flat versus frequencies.

4. PROPOSED RESEARCH

In the future, we propose to study the effect of Landau damping and particle collisions in detail. Experimentally, we will measure the ion acoustic waves in a variety of different situations including different neutral pressures and plasma densities.

REFERENCES

- [1]. B. Basu and J. R. Jasperse, Marginal stability of ion-acoustic waves in a weakly collisional two-temperature plasma without a current, *Phys. Rev.*, 38, 3785, 1988.

Table 1. Experimental and theoretical wave parameters

	experiment	theory
wave velocity (km/s)	1.86	1.90
Landau damping rate (cm^{-1})	$0.0025 \times f(\text{kHz})$	$0.0024 \times f(\text{kHz})$
collisional damping rate (cm^{-1})	-0.0735	i-n: 0.002; e-i: -0.068 total: -0.066




Figure 1. Schematic of the plasma chamber for measuring ion acoustic waves.




Figure 2. The I-V curve taken from Langmuir probe




Figure 3. The waves measured in different frequencies




Figure 3. (cont.)





Figure 4. Normalized wave number and damping rate versus ratio of ion to electron temperatures. The solid lines are from Equation (1a) and (1b) and the dashed lines are from Equation (2) ($\omega/\omega_{pi}=0.1$).






Figure 5. The wave number and damping rate versus frequency.



(a)



(b)

Figure 6. (a). Wave number versus frequency; (b). Damping rate versus frequency